


Optimal Controlled Phasegates for Ultracold Atoms in an Optical Lattice at the Quantum Speed Limit



Michael Goerz¹, Tommaso Calarco², Christiane P. Koch^{1,3}

1: Institut für Theoretische Physik, FU Berlin, 2: Institut für Quanteninformationsverarbeitung, Universität Ulm, 3: Institut für Physik, Universität Kassel

Summary

We study controlled phasegates for ultracold atoms in an optical lattice [1]. The qubits are encoded in the electronic states. A shaped laser pulse drives transitions between the ground and electronically excited states where the atoms are subject to a long-range $1/R^3$ interaction. We fully account for this interaction and use optimal control theory to calculate the pulses. This allows us to determine the minimum pulse duration, respectively the gate time T that is required to obtain high fidelity. We find the gate time to be limited either by the interaction strength in the excited state or by the ground state vibrational motion in the trap. The latter needs to be resolved in order to fully restore the motional state of the atoms at the end of the gate.

Universal Quantum Computing

The set of all one-qubit gates plus the two-qubit CNOT is universal. More generally, the CNOT is equivalent to the controlled phasegate, combined with Hadamard and X-gates.

$$\hat{O}(\phi) = \begin{pmatrix} e^{i\phi} & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}; \quad CNOT = \begin{array}{c} \text{---} \text{X} \text{---} \\ | \\ \text{---} \text{H} \text{---} \text{X} \text{---} \text{O}(\pi) \text{---} \text{X} \text{---} \text{H} \text{---} \end{array}$$

Qubit Encoding in Calcium

$$\hat{H}_{2q} = \begin{pmatrix} E_0 & 0 & \mu\epsilon(t) \\ 0 & E_1 & 0 \\ \mu\epsilon(t) & 0 & E_a \end{pmatrix} \otimes \mathbb{1}_{1q} \otimes \mathbb{1}_{x_1} + \mathbb{1}_{1q} \otimes \begin{pmatrix} E_0 & 0 & \mu\epsilon(t) \\ 0 & E_1 & 0 \\ \mu\epsilon(t) & 0 & E_a \end{pmatrix} \otimes \mathbb{1}_{x_2} + \sum_{ij} \hat{V}_{BO}^{(ij)}(|x_2 - x_1\rangle) + \sum_{ij} \hat{V}_{\text{trap}}^{(ij)}(x_1, x_2)$$


initialize to trap ground state $\Psi_{00,i}(R) \approx \left(\frac{\mu\omega_0}{4\pi\hbar}\right)^{1/4} \sum_{\pm} e^{-\frac{m\omega_0}{2\hbar}(d \pm R)^2}$

- Parameters:
- Use actual calcium $B^1\Sigma_u^+$ interaction potential at $d = 5$ nm.
 - Use generic dipole-dipole interaction with variable C_3 at $d = 200$ nm.

$$\begin{array}{c} |aa\rangle \\ |0a\rangle \\ |a0\rangle \\ |00\rangle \end{array} = -\frac{C_3}{R^3}$$

OCT: Finding an Optimized Pulse

Change pulse iteratively to minimize [2, 3] $J = -F + \int \frac{\alpha}{S(t)} \Delta\epsilon(t) dt$, $F = \frac{1}{N} \Re[\text{tr}(\hat{O}^\dagger \hat{U})]$



$$\Delta\epsilon(t) = \frac{S(t)}{\alpha} \Im \left[\sum_{k=1}^N \frac{1}{2} \langle \Psi_{ik} | \hat{O}^\dagger \hat{U}^\dagger(T \rightarrow t, \epsilon^{(0)}) \hat{\mu} \hat{U}(0 \rightarrow t, \epsilon^{(1)}) | \Psi_{ik} \rangle \right]$$


One-Qubit and Two-Qubit Phases

	full optimization scheme	reduced optimization scheme
system phases	$\phi_{00}, \phi_{01}, \phi_{10}, \phi_{11}$	$\phi_{00}, \phi_0, \phi_1$
optimization targets	$ 00\rangle \rightarrow e^{i(\phi+\phi_T)} 00\rangle$ $ 01\rangle \rightarrow e^{i\phi_T} 01\rangle$ $ 10\rangle \rightarrow e^{i\phi_T} 10\rangle$ $ 11\rangle \rightarrow e^{i\phi_T} 11\rangle$	$ 00\rangle \rightarrow e^{i(\phi+\phi_T)} 00\rangle$ $ 0\rangle \rightarrow e^{i\phi_T/2} 0\rangle$ $ 1\rangle \rightarrow e^{i\phi_T/2} 1\rangle$
gate phases	ϕ_{00} $\phi_{10} = \phi_{01}$ ϕ_{11}	$= \phi_{00}$ $= \phi_0 + \phi_1$ $= 2\phi_1$
true two-qubit phase	$\chi = \phi_{00} - \phi_{01} - \phi_{10} + \phi_{11}$	$\chi = \phi_{00} - 2\phi_0$


True two-qubit phase χ from Cartan decomposition [4].

$$\hat{U} = \hat{U}_1 \hat{O}(\chi) \hat{U}_2,$$


where \hat{U}_1 and \hat{U}_2 are purely local operations.






Results



Phase Dynamics for $T = 1.23$ ps in two-qubit and single-qubit system



Phase Dynamics for $T = 290$ ps in two-qubit and single-qubit system



References

- [1] T. Calarco et al., *Phys. Rev. A* **61**, 022304 (2004)
- [2] J. P. Palao, R. Kosloff, *Phys. Rev. Lett.* **89**, 188301 (2002), *Phys. Rev. A* **68**, 062308 (2003).
- [3] C. P. Koch et. al. *Phys. Rev. A* **70**, 013402 (2004).
- [4] J. Zhang et. al. *Phys. Rev. A.* **67**, 042313 (2003).