

# Chebychev Propagator for Inhomogeneous Schrödinger Equations

Michael Goerz

May 16, 2011

## Solving the Schrödinger Equation

### Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle; \quad \text{e.g. } \hat{H} = \begin{pmatrix} V_1(R) & \mu\epsilon(t) \\ \mu\epsilon(t) & V_2(R) \end{pmatrix}$$

## Solving the Schrödinger Equation

### Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle; \quad \text{e.g. } \hat{H} = \begin{pmatrix} V_1(R) & \mu\epsilon(t) \\ \mu\epsilon(t) & V_2(R) \end{pmatrix}$$

### Solution

$$|\Psi(t)\rangle = e^{-i\hat{H}t} |\Psi_0\rangle \quad \text{if } \hat{H} \text{ not time dependent}$$

## Solving the Schrödinger Equation

### Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle; \quad \text{e.g. } \hat{H} = \begin{pmatrix} V_1(R) & \mu\epsilon(t) \\ \mu\epsilon(t) & V_2(R) \end{pmatrix}$$

### Solution

$$|\Psi(t + \Delta t)\rangle = e^{-i\hat{H}\Delta t} |\Psi(t)\rangle \rightarrow \text{piecewise constant pulses}$$

## Solving the Schrödinger Equation

### Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle; \quad \text{e.g. } \hat{H} = \begin{pmatrix} V_1(R) & \mu\epsilon(t) \\ \mu\epsilon(t) & V_2(R) \end{pmatrix}$$

### Solution

$$|\Psi(t + \Delta t)\rangle = e^{-i\hat{H}\Delta t} |\Psi(t)\rangle \rightarrow \text{piecewise constant pulses}$$

### Evaluation of the Time Evolution Operator

$$\text{Expand into series: } e^{-i\hat{H}t} \rightarrow \sum_{k=1}^N a_k P_k(\hat{H})$$

## Solving the Schrödinger Equation

### Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle; \quad \text{e.g. } \hat{H} = \begin{pmatrix} V_1(R) & \mu\epsilon(t) \\ \mu\epsilon(t) & V_2(R) \end{pmatrix}$$

### Solution

$$|\Psi(t + \Delta t)\rangle = e^{-i\hat{H}\Delta t} |\Psi(t)\rangle \rightarrow \text{piecewise constant pulses}$$

### Evaluation of the Time Evolution Operator

$$\text{Expand into series: } e^{-i\hat{H}t} \rightarrow \sum_{k=1}^N a_k P_k(\hat{H})$$

cf. Runge-Kutta: solving the differential equation, instead of evaluating the analytical solution

## Chebyshev Polynomials

### Properties of Chebyshev polynomials

- $P_0 = 1$ ,  $P_1(x) = x$ ,  $P_n(x) = 2xP_{n-1}(x) - P_{n-2}(x)$
- Defined over range  $[-1, 1] \rightarrow$  normalize Hamiltonian

$$\hat{H}_{\text{norm}} = 2 \frac{\hat{H} - E_{\min} \mathbb{1}}{\Delta E} - \mathbb{1}$$

- Fastest converging polynomial expansion
- $P_n(x) = \cos(n\theta)$  with  $\theta = \arccos(x) \rightarrow$  Cosine transform for coefficients

### Chebyshev coefficients

- Expansion coefficients  $a_n$  for function  $f(x)$ :

$$a_n = \frac{2 - \delta_{n0}}{\pi} \int_{-1}^{+1} \frac{f(x)P_n(x)}{\sqrt{1-x^2}} dx$$

- For  $f(\hat{H}_{\text{norm}}) = e^{-i\hat{H}_{\text{norm}}t}$ :  $a_n \rightarrow$  Bessel functions.

## Inhomogeneous Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + |\Phi(t)\rangle$$




## Inhomogeneous Schrödinger Equation

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + |\Phi(t)\rangle$$

Note: not the same as *nonlinear* SE:

$$\text{e.g. } \frac{\partial}{\partial t} |\Psi(t)\rangle = (\hat{H} + |\Psi(t)|^2) |\Psi(t)\rangle$$


## OCT with State-Dependent Costs



### Optimization Functional

$$J = -F[\Psi_t] + \int g_a[\epsilon(t)] dt + \int g_b[\Psi(t)] dt; \quad g_b[\Psi] = \lambda_b \langle \Psi(t) | \hat{P}_{\text{allow}} | \Psi(t) \rangle$$

## OCT with State-Dependent Costs




### Optimization Functional

$$J = -F[\Psi_t] + \int g_a[\epsilon(t)] dt + \int g_b[\Psi(t)] dt; \quad g_b[\Psi] = \lambda_b \langle \Psi(t) | \hat{P}_{\text{allow}} | \Psi(t) \rangle$$

also: time-dependent targets

## Reminder: Krotov



Pulse update by matching forward- and backward-propagated states

## Inhomogeneous Backward-Propagation

Daniel: Second Order Krotov Preprint (arXiv:1008.5126v1)

### Optimization Functional

$$J = -F[\Psi_t] + \int g_a[\epsilon(t)] dt + \int g_b[\Psi(t)] dt; \quad g_b[\Psi] = \lambda_b \langle \Psi(t) | \hat{\mathbf{P}}_{\text{allow}} | \Psi(t) \rangle$$

### Pulse Update

$$\Delta\epsilon \propto -\Im \langle \chi^{(0)}(t) | \hat{\mu} | \phi^{(1)}(t) \rangle$$

### Backward Propagation

$$\begin{aligned} \frac{d}{dt} |\chi^{(0)}(t)\rangle &= -\frac{i}{\hbar} \hat{\mathbf{H}}^\dagger[\varphi^{(0)}, \epsilon^{(0)}] |\chi^{(0)}(t)\rangle + \nabla_{\langle \varphi | g_b | \varphi^{(0)}(t) \rangle} \\ |\chi^{(0)}(T)\rangle &= -\nabla_{\langle \varphi | J_T | \varphi^{(0)}(T) \rangle} \end{aligned}$$

## Inhomogeneous Backward-Propagation

Daniel: Second Order Krotov Preprint (arXiv:1008.5126v1)

### Optimization Functional

$$J = -F[\Psi_t] + \int g_a[\epsilon(t)] dt + \int g_b[\Psi(t)] dt; \quad g_b[\Psi] = \lambda_b \langle \Psi(t) | \hat{\mathbf{P}}_{\text{allow}} | \Psi(t) \rangle$$

### Pulse Update

$$\Delta\epsilon \propto -\Im \langle \chi^{(0)}(t) | \hat{\mu} | \phi^{(1)}(t) \rangle$$

### Backward Propagation

$$\begin{aligned} \frac{d}{dt} |\chi^{(0)}(t)\rangle &= -\frac{i}{\hbar} \hat{\mathbf{H}}^\dagger [\varphi^{(0)}, \epsilon^{(0)}] |\chi^{(0)}(t)\rangle + \hat{\mathbf{P}}_{\text{allow}} |\varphi^{(0)}(t)\rangle \\ |\chi^{(0)}(T)\rangle &= -\nabla_{\langle \varphi | J_T | \varphi^{(0)}(T) \rangle} \end{aligned}$$

# Solving the Inhomogeneous Schrödinger Equation Numerically

## References

### [1] JCP 130, 124108 (2009)

THE JOURNAL OF CHEMICAL PHYSICS **130**, 124108 (2009)

#### A Chebyshev propagator for inhomogeneous Schrödinger equations

Mamadou Ndong,<sup>1</sup> Hillel Tal-Ezer,<sup>2</sup> Ronnie Kosloff,<sup>3</sup> and Christiane P. Koch<sup>1,a)</sup>

<sup>1</sup>*Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany*

<sup>2</sup>*School of Computer Sciences, The Academic College of Tel-Aviv Yaffo, 2 Rabenu Yeruham St., Tel-Aviv 61803, Israel*

<sup>3</sup>*Department of Physical Chemistry and The Fritz Haber Research Center, The Hebrew University, Jerusalem 91904, Israel*

### [2] JCP 132, 064105 (2010)

THE JOURNAL OF CHEMICAL PHYSICS **132**, 064105 (2010)

#### A Chebyshev propagator with iterative time ordering for explicitly time-dependent Hamiltonians

Mamadou Ndong,<sup>1</sup> Hillel Tal-Ezer,<sup>2</sup> Ronnie Kosloff,<sup>3</sup> and Christiane P. Koch<sup>1,a)</sup>

<sup>1</sup>*Institut für Theoretische Physik, Freie Universität Berlin, Arnimallee 14, 14195 Berlin, Germany*

<sup>2</sup>*School of Computer Sciences, The Academic College of Tel Aviv-Yaffo, Rabenu Yeruham St., Tel-Aviv 61803, Israel*

<sup>3</sup>*Institute of Chemistry and The Fritz Haber Research Center, The Hebrew University, Jerusalem 91904, Israel*



## Treating the Inhomogeneity in Order $m$

### Inhomogeneous SE

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + |\Phi(t)\rangle$$

### Expansion of $\Phi(t)$

$$|\Phi(t)\rangle_m = \sum_{j=0}^{m-1} |\bar{\Phi}_j\rangle P_j(\bar{t})$$

- Expand inhomogeneous term in Chebychev series

## Treating the Inhomogeneity in Order $m$

### Inhomogeneous SE

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + |\Phi(t)\rangle$$

### Expansion of $\Phi(t)$

$$|\Phi(t)\rangle_m = \sum_{j=0}^{m-1} |\bar{\Phi}_j\rangle P_j(\bar{t}) \equiv \sum_{j=0}^{m-1} \frac{t^j}{j!} |\Phi^{(j)}\rangle$$

- Expand inhomogeneous term in Chebychev series
- Reorder into power series (or use Taylor to begin with)

## Treating the Inhomogeneity in Order $m$

### Inhomogeneous SE

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + |\Phi(t)\rangle$$

### Expansion of $\Phi(t)$

$$|\Phi(t)\rangle_m = \sum_{j=0}^{m-1} |\bar{\Phi}_j\rangle P_j(\bar{t}) \equiv \sum_{j=0}^{m-1} \frac{t^j}{j!} |\Phi^{(j)}\rangle$$

- Expand inhomogeneous term in Chebychev series
- Reorder into power series (or use Taylor to begin with)
- Decide on which order to solve: 1, 2, 3, maybe 4
- The smaller the order, the smaller  $\Delta t$  has to be

## The Analytical Solution

### Inhomogeneous SE ( $\Phi$ to order $m$ )

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + \sum_{j=0}^{m-1} \frac{t^j}{j!} |\Phi^{(j)}\rangle$$

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle$$

$$f_m = (-i\hat{H})^{-m} \left( e^{-i\hat{H}t} - \sum_{j=0}^{m-1} \frac{(-i\hat{H}t)^j}{j!} \right) \quad \lambda^{(0)} = |\Psi_0\rangle$$

$$\lambda^{(j)} = -i\hat{H} |\lambda^{(j-1)}\rangle + |\Phi^{(j-1)}\rangle$$

## The Analytical Solution

### Inhomogeneous SE ( $\Phi$ to order $m$ )

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + \sum_{j=0}^{m-1} \frac{t^j}{j!} |\Phi^{(j)}\rangle$$

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle$$

$$f_m = (-i\hat{H})^{-m} \left( e^{-i\hat{H}t} - \sum_{j=0}^{m-1} \frac{(-i\hat{H}t)^j}{j!} \right) \quad \begin{aligned} \lambda^{(0)} &= |\Psi_0\rangle \\ \lambda^{(j)} &= -i\hat{H} |\lambda^{(j-1)}\rangle + |\Phi^{(j-1)}\rangle \end{aligned}$$

e.g.  $|\Psi(t)\rangle_{(3)} = |\Psi_0\rangle + t |\lambda^{(1)}\rangle + \frac{t^2}{2} |\lambda^{(2)}\rangle + f_3(\hat{H}) |\lambda^{(3)}\rangle$ , with

$$f_3(\hat{H}) = (-i\hat{H})^{-3} \left( e^{-i\hat{H}t} - 1 - i\hat{H}t \right)$$

## The Analytical Solution

### Inhomogeneous SE ( $\Phi$ to order $m$ )

$$\frac{\partial}{\partial t} |\Psi(t)\rangle = \hat{H} |\Psi(t)\rangle + \sum_{j=0}^{m-1} \frac{t^j}{j!} |\Phi^{(j)}\rangle$$

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle$$

$$f_m = (-i\hat{H})^{-m} \left( e^{-i\hat{H}t} - \sum_{j=0}^{m-1} \frac{(-i\hat{H}t)^j}{j!} \right) \quad \lambda^{(0)} = |\Psi_0\rangle$$

$$\lambda^{(j)} = -i\hat{H} |\lambda^{(j-1)}\rangle + |\Phi^{(j-1)}\rangle$$

e.g.  $|\Psi(t)\rangle_{(0)} = e^{-i\hat{H}t} |\Psi_0\rangle \rightarrow$  homogeneous propagation

## Chebychev Propagation

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle; \quad |\lambda\rangle \sim \{|\Phi^{(j)}\rangle\}$$

### Idea

Evaluate  $f_m(\hat{H})$  by expanding it into Chebychev Polynomials  
(just like for the “standard” Chebychev propagator with  $f_0(\hat{H}) = e^{i\hat{H}t}$ )

## Chebyshev Propagation

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle; \quad |\lambda\rangle \sim \{|\Phi^{(j)}\rangle\}$$

### Idea

Evaluate  $f_m(\hat{H})$  by expanding it into Chebyshev Polynomials  
(just like for the “standard” Chebyshev propagator with  $f_0(\hat{H}) = e^{i\hat{H}t}$ )

### Algorithm Outline (for fixed $m$ )

For each time step:

- determine  $\{|\Phi^{(j)}\rangle\}$  and from that  $\{\lambda^{(j)}\}$
- run through the Chebyshev series for  $f_m$
- sum everything up, yielding  $|\Psi(t)\rangle_{(m)}$



# Details of the Algorithm

## Global Initialization (before any actual propagation)

### Calculate Chebychev Coefficients

Calculate the Chebychev expansion coefficients  $a_n$  for  $f_m(\hat{H})$ , for the chosen order  $m$ .

## Global Initialization (before any actual propagation)

### Calculate Chebychev Coefficients

Calculate the Chebychev expansion coefficients  $a_n$  for  $f_m(\hat{\mathbf{H}})$ , for the chosen order  $m$ .

- $a_n$  *cannot* be calculated analytically (like for the standard Chebychev)
- Calculation of  $a_n$  is done through a fast cosine-transform:

$$a_n = \frac{2 - \delta_{n0}}{N} \sum_{k=0}^{N-1} f_m(\theta_k) \cos(n\theta_k)$$

- $\hat{\mathbf{H}}$  needs to be normalized  $\rightarrow a_n$  might have to be re-calculated if spectral radius changes (after each OCT iteration)
- On a non-equidistant time grid, the  $a_n$  would have to be re-calculated at every time step
- *For small  $\hat{\mathbf{H}}$ , the term  $(-i\hat{\mathbf{H}})^{-m}$  might lead to numerical instability. We could use Taylor instead. ... ?*

## Local Initialization (at every time step)

### Calculate Expansion of Inhomogeneous Term

Calculate all necessary  $|\Phi^{(j)}\rangle$  (i.e. up to order  $m$ ) to approximate the local  $\Phi(t_i)$ , either by an intermediate Chebychev expansion, followed by calculation of coefficients in the power series, or by a direct Taylor series.

## Local Initialization (at every time step)

### Calculate Expansion of Inhomogeneous Term

Calculate all necessary  $|\Phi^{(j)}\rangle$  (i.e. up to order  $m$ ) to approximate the local  $\Phi(t_i)$ , either by an intermediate Chebychev expansion, followed by calculation of coefficients in the power series, or by a direct Taylor series.

Calculation via Taylor:

- Calculate derivatives through FFT

Calculation via Chebychev:

- Sample  $\Phi(t)$  at intermediate points around  $t$  by interpolation (splining should be fine)
- Calculate Chebychev coefficients  $|\bar{\Phi}_j\rangle$  by cosine transform
- Calculate  $|\Phi^{(j)}\rangle$  from  $|\bar{\Phi}_j\rangle$  by formulas in References (just collect the powers)

## Propagation Step

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle$$

- Calculate  $|\lambda^{(j)}\rangle, j = 0 \dots m-1$

$$\lambda^{(j)} = -i\hat{H} |\lambda^{(j-1)}\rangle + |\phi^{(j-1)}\rangle$$

- Calculate  $f_m(\hat{H}) |\lambda^{(m)}\rangle$  by Chebychev recursion (Choose  $N$  to reach machine precision)

$$f_m(\hat{H}) |\lambda^{(m)}\rangle = \sum_{n=1}^N a_n P_n(\hat{H}) |\lambda^{(m)}\rangle$$

$$P_n(\hat{H}) = 2\hat{H}P_{n-1}(\hat{H}) - P_{n-2}(\hat{H})$$

- Calculate  $|\Psi(t)\rangle_{(m)}$

# Inhomogeneous Chebychev in QDYN

## Propagation Routines (prop.f90)

### prop

```
subroutine prop(psi, grid, ham, work, para, from_ti, to_ti, bw, &  
& info_hook, pulses, alt_pulse, upd_hook, storage)
```

### prop\_step

```
subroutine prop_step(psi, grid, ham, work, para, ti, bw, &  
& pulses, alt_pulses, alt_pulse, upd_hook)
```



## Propagation Routines (prop.f90)

### prop

```
subroutine prop(psi, grid, ham, work, para, from_ti, to_ti, bw, &  
& info_hook, pulses, alt_pulse, upd_hook, storage)
```

### prop with optional inhomogeneous term

```
subroutine prop(psi, grid, ham, work, para, from_ti, to_ti, bw, inh_psi, &  
& get_inh_phi, info_hook, pulses, alt_pulse, upd_hook, storage)
```

- `inh_psi`: stored array of forward-propagated states
- `get_inh_phi`: function that calculates  $|\Phi\rangle$  from  $|\Psi\rangle$  (e.g.  $\hat{P}_{\text{allow}}|\Psi\rangle$ ).

### prop\_step

```
subroutine prop_step(psi, grid, ham, work, para, ti, bw, &  
& pulses, alt_pulses, alt_pulse, upd_hook)
```

### prop\_step with optional inhomogeneous term

```
subroutine prop_step(psi, grid, ham, work, para, ti, bw, inh_psi, &  
& get_inh_phi, pulses, alt_pulses, alt_pulse, upd_hook)
```

## Calculation of Chebychev Coefficients

### Calculate Chebychev Coefficients

Calculate the Chebychev expansion coefficients  $a_n$  for  $f_m(\hat{H})$ , for the chosen order  $m$ .

in `inhom_cheby.f90`:

subroutine `init_inh_cheby`(ham, wcheby, order, para)

- Use the same work array (`wcheby`) as for the homogeneous Chebychev propagation.
- Sufficient  $a_n$  are calculated and stored in `wcheby` to reach machine precision
- Watch out for numerical instability ( $\rightarrow$  Taylor)

## Expansion of Inhomogeneous Term

### Calculate Expansion of Inhomogeneous Term

Calculate all necessary  $|\Phi^{(j)}\rangle$  (i.e. up to order  $m$ ) to approximate the local  $\Phi(t_i)$ , either by an intermediate Chebychev expansion, followed by calculation of coefficients in the power series, or by a direct Taylor series.

in `inhom_cheby.f90`:

```
subroutine expand_inh_phi(inh_psi, get_inh_phi, order, phi_coeffs)
```

- `phi_coeffs` stores the  $|\Phi^{(j)}\rangle$  (power series)
- invoke `monic_transf` to calculate  $|\Phi^{(j)}\rangle$  from  $|\bar{\Phi}_j\rangle$

Continue with calculation of  $|\lambda^{(j)}\rangle$ :

```
subroutine get_inh_lambda(lambda, phi, ham, ...)
```

## Performing the Propagation Step

### Solution

$$|\Psi(t)\rangle_{(m)} = \sum_{j=0}^{m-1} \frac{t^j}{j!} |\lambda^{(j)}\rangle + f_m(\hat{H}) |\lambda^{(m)}\rangle$$

in `inhom_cheby.f90`:

```
subroutine inhom_cheby(psi, work, ham, grid, lambda, para, dt, ti, &  
& alt_pulses, alt_pulse)
```

- Identical interface to `cheby`, except for  $\lambda$
- Inhomogeneous Chebychev coefficients are implicit in `work`